

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 7/06, 7/18, 14/00, 19/00, 99/18, A61K 38/00, 38/08		A1	(11) International Publication Number: WO 00/11022
			(43) International Publication Date: 2 March 2000 (02.03.00)
(21) International Application Number: PCT/US99/19381		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 20 August 1999 (20.08.99)			
(30) Priority Data:			
60/097,210	20 August 1998 (20.08.98)	US	
60/141,169	25 June 1999 (25.06.99)	US	
09/378,019	19 August 1999 (19.08.99)	US	
(71)(72) Applicants and Inventors: STEWART, John, M. [US/US]; 3690 East Dartmouth Avenue, Denver, CO 80210 (US). CHAN, Daniel, C., F. [US/US]; 3691 South Quebec, Denver, CO 80237 (US). GERA, Lojos [HU/US]; 880 Cherry Street #208, Denver, CO 80220 (US). YORK, Eunice [US/US]; 3102 South Gaylord Street, Englewood, CO 80110 (US). BUNN, Paul [US/US]; 630 Sundown Lane, Evergreen, CO 80439 (US).		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(74) Agent: JOHNSON, Kristine, H.; Macheledt Bales & Johnson LLP, The Opera Galleria, Suite 219, 123 North College Avenue, Fort Collins, CO 80524 (US).			
(54) Title: ANTI-CANCER COMPOUNDS AND METHODS RELATED THERETO			
(57) Abstract			
<p>The present invention provides compounds useful to inhibit tumor growth and to induce apoptosis. In general, the anti-cancer agents (ACA) are described by the formula (I): [ACA]_n-X wherein, X is a linker group having 2-5 functional groups or is absent, n = 1, and ACA is selected from the group consisting of Formula (II), Formula (III), Formula (IV), Formula (V), and Formula (VI), as described herein. Other compounds described herein are defined by the Formula (VII), as described herein.</p>			
BEST AVAILABLE COPY			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

5 **ANTI-CANCER COMPOUNDS AND METHODS RELATED THERETO**

The present application claims priority to US Provisional Patent application serial Number 60/097,210, filed August 20, 1998 and US Provisional Patent application serial number 60/141,169, filed June 25, 1999.

10

Field of the Invention

The present invention relates generally to the field of cancer treatments, as well as to the field of peptide and non-peptide pharmaceutical compounds.

15

BACKGROUND OF THE INVENTION

Many lung and prostate cancers, of which small cell lung cancer (SCLC) is a prime example, have a neuroendocrine phenotype, and their growth is stimulated by
20 neuropeptides. Antagonists of several peptides (e.g. bradykinin, substance P, bombesin) have been used in experimental treatment of models of SCLC in animals. Among the most potent of the peptides examined thus far, crosslinked dimers of certain bradykinin antagonist peptides have been efficacious both *in vitro* and *in vivo* against strains of SCLC and other tumors (Chan et al., *Immunopharmacology* 33: 201-204, 1996; Stewart et al.,
25 *Can. J. Physiol. Pharmacol.* 75: 719-724, 1997; Stewart et al., US Patent Application 5,849,863, issued 12/15/98). Prostate cancers show a similar neuroendocrine phenotype and are susceptible to neuropeptide antagonists.

SUMMARY OF THE INVENTION

30


The present invention provides anti-cancer agents (ACA) comprised of a range of novel amino acid derivatives and small peptides having the ability to inhibit growth of SCLC and certain other tumor cell lines (such as non-small cell lung cancer (NSCLC) and prostate cancer) in standard *in vitro* tests, as well as certain monomeric peptides that show
35 inhibition of tumor growth *in vivo*. Certain of the peptides have a general structural relationship to carboxy-terminal fragments of bradykinin antagonists, but the non-peptides show no such general relationship. Monomers, dimers, trimers, tetramers, pentamers and

5 cyclized analogs of the novel molecules are described. The new compounds are tested for bradykinin antagonist activity in standard assays, but there is no apparent relationship between bradykinin antagonist activity and cytolytic potency. All of the molecules described possess both hydrophobic (usually aromatic) and basic groups in their structures. Without being held to one particular theory, it appears that the compounds
10 function by stimulation of cell death (apoptosis) in the tumor cells.

The present invention also provides compounds and methods for inhibiting cancer by administering to a subject afflicted with cancer (ie. of the lung or prostate) a therapeutically effective amount of one or more of the compounds herein described.

In general, the anti-cancer compounds are described by the formula:

15



wherein X is a linker having 2-5 functional groups or is absent, $n = 1-5$, and ACA is selected from the group consisting of Formula II, Formula III, Formula IV, Formula V, and Formula VI. Other compounds described herein are defined by the Formula VII. The
20 specifics regarding structure are enumerated in the Detailed Description, Examples and Claims. Certain physical characteristics are enumerated in the Examples as well as the Detailed Description, Examples and Claims.

BRIEF DESCRIPTION OF THE DRAWINGS

25 Figure 1 shows inhibition of growth *in vivo* of SCLC strain SHP-77 by B10054.
Figure 2 shows inhibition of growth *in vivo* of NSCLC strain A-549 by M620.
Figure 3 shows inhibition of growth *in vivo* of SCLC strain SHP-77 by B9430.
Figure 4 shows inhibition of growth *in vivo* of SCLC strain SHP-77 by B10238
Figure 5 shows inhibition of growth *in vivo* of SCLC strain SHP-77 by M570, both
30 as the trifluoroacetate salt and as the hydrochloride salt.
Figure 6 shows inhibition of growth *in vivo* of SCLC strain SHP-77 by M822.
Figure 7 shows inhibition of growth *in vivo* of SCLC strain SHP-77 by M638.

5

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a range of monomeric, dimeric, trimeric, tetrameric, pentameric and cyclic small peptides and peptide mimics that are effective as anti-cancer agents.

In general, the anti-cancer agents (ACA) are described by the formula:

10



wherein X is a linker group having 2-5 functional groups or is absent, $n = 1$, and ACA is selected from the group consisting of Formula II, Formula III, Formula IV, Formula V, and Formula VI, as described herein. Other compounds described herein are defined by the
15 Formula VII, as described herein.

X can be any linking group which does not interfere with the inhibitory activity of the monomer-linker or oligomerized product using *bis*-imido-esters, *bis*-maleimidoalkanes such as *bis*-maleimidohexane, dicarboxylic acids, tricarboxylic acids, tetracarboxylic acids and multi carboxylic acids. Alkane groups may be substituted with alkyl, amino, carboxyl,
20 halogen, hydroxy, mercapto or methoxy groups. Aminoalkyl, aromatic or cycloalkyl polycarboxylic acids, heterocyclic polycarboxylic acids, carboxylic anhydrides and polyoxyethylene linkers may also be used. For C-terminal crosslinking, X may be any diamino or polyamino alkane, cycloalkane, aromatic, heterocyclic diamine, polyamine or other substituted chelating agent (for example: diethylenetriaminepentaacetic dianhydride,
25 ethylenediaminetetraacetic dianhydride, etc.). Polyamino-polycarboxylic acids may also be used to make heteromers (such as ethylenediamine-N,N'-diacetic acid, etc.).

The linkage may be at the N-terminal or the C-terminal or at any position of the ACA sequence through side-chain functional groups. The linker may have any chain length.

30

For dimers, there is a correspondence between linker length and cytotoxicity. Alkyl chains of 8 carbons or more are preferred, with those of 8 to 18 carbons being most preferred. Examples of preferred dimer linkers for the α -amino at the N-terminal or for a basic side-chain group at any position of ACA include ADA, BTAC, DDD, DDS, DTP,


- 5 EGS, EOPC, HDD, HFG, PFS, SBEC, SUB, SUIM and TDIM. For dimerization through the C-terminal carboxyl or any side-chain carboxyl in ACA, the preferred linkers include DDA, DEA, EDA, EDP and HAD. Any di-functional molecule can be used.

For trimers, linkers for basic groups include BTAC, BTC, CHTC, CTAC and TREN-(Suc)₃; for carboxyl groups, TREN. Any tri-functional molecule can be used.

- 10 For tetramers, linkers can be BAPTA, CPTA, EDTA, EGTA, ETТА, or any tetra-functional molecule.

For pentamers, the linker can be DTPA or any pentameric functional molecule. Compounds formed by ACA and a linker X may be homo or hetero multimers.

- 15 [Formula II] comprises:



- 20 wherein R, A, B, C, D, E, F, G, H, I, J, and K are selected from the following or may be absent, and wherein K is Arg or an Arg derivative:

R	A -1	B 0	C 1	D 2	E 3	F 4	G 5	H 6	I 7	J 8	K 9
Absent or 3,3DP Aaa Ac	Absent or DmK Lys Lys(εF lu)	Absent or Apc Arg DArg	Absent or ApC Arg DmK	Absent Or MeP Nig NMF	Absent or Hyp Pro	Absent or Ava BAIa Dpr	Absent or Add Aud CpG	Absent or Arg Gly Pac	Absent or 2Nal DCpG DF5F	Absent or 2Nal 2Nal- NH ₂	Absent or Arg Arg(H) Arg- CH ₂ O H Arg- NH ₂ Arg(N O ₂) Arg- OMe DArg DArg- NH ₂ DArg(NO ₂)
Aca	NiK	DLys	NiK	Pro		Eac	DDMF	Pac	Dlgl	3,4F2F	
BApp	PzO	DmK	NiO			Gly	DMF	Ser	DPFF	3Pal	
Cca		DniK	PaF				Eac	Thr	DPhe	Ac6c	
Cin Dca		DpaF DPzK	PzO				Igl Lys		DTic Gly	Aic Ana	
Dcg		DPzO					Pac		mABz	Apb	
Dhq Dmac Dpa		Lys NiK PaF					Phe Thi		pABz Pac PaF(Dc g)	Apb Atpc Bip	
F5bz F5c		PzO DArg- (NO ₂)	Arg- (NO ₂)						pAmb	Cmp CpG	

F5pa
Gun

Hxa

Mca
Mcg

Moti
Pcc
Ppa
Pya


Saa
Ste

Tfmc

DhPhe
Dpr(Fb
z)
Dpr(Pa
a)
F5F
F5F-
NH₂
Hphe
Ica
Igl
Igl-
NH₂
Ileu
Lys(C
H₃)₃
Lys(F5
bz)
Mapa
MBC
MFF
Nc6G
Nc7G
NMF
OBS
OBT
OBY
OC2Y
Oic
Oic-
NH₂
PABz
Pac
PaF(F5
c)
PaF(Fb
z)
PaF(M
cg)
PaF(Pp
a)
PaF(Si
n)
pAmb
pAPa
PCF
PdF
PFF
PFF-
NH₂
Phe
PNF
Thi
Tic
Trp
Trx
Tyr

5

[Formula III] comprises:




- 10 wherein R, A, B, C, D, E, and F are selected from the following or may be absent, and wherein F is not Arg or an Arg derivative:

R	A 1	B 2	C 3	D 4	E 5	F 6
Absent or 2,2Dp 3,3Dp	Absent or DArg DArg(NO ₂))	Absent or Arg	Absent or Add Aud	Absent or 2Nal 3Pal	Absent or 1Nal 2Nal	Absent or 2Nal 3Pal
Aaa			Ava	Arg	2Nap	ABza
Ac			Eac	Arg(Tos)	3Pal	ABza
Aca			Lys	Atcp	Apa	Ama
Boc			Pac	D2Nal	Arg	Ampy
Chc				DArg	Arg-NH ₂	Ampz
Cin				DArg(Tos)	Asp	Apa
Ctim				DF5F	Atc	Api
Dca				DIgl	Atcp	Aptp
Dcg				DPFF	Bip	Aqd
Dhq				Eac	BtA	Aqu
Dmac				F5F	Cys(Meb)	Arg(H)
Dns				Gly	Cys(SO ₃ H)	Arg- CH ₂ OH
Dpa				His	D2Nal	Arg-NH ₂
F5c				Igl	DArg	Arg-OMe
F5pa				mABz	DArg-NH ₂	Asp
F5po				OC2Y	F5F	Asp(Aqu)
Gbc				Pac	Glu	Atcp
Gun				PFF	Gly	Atmp
Hxa					Igl	AtmpO
Mcg					Inp	Atpm
Mse					Iqa	Cyh
Pya					mABz	Dmab
Seb					MC2Y	Dmm
Sin					N-Dmb- Tyr(Bz)- OMe	Dmp
Sul					OC2Y	Dpea
Tfmc					OCIY	Dpma
Tha					Oic	Dpr(Dcg-2- Nap)
					pABz	Ecap
					PaF(Mes)	F5F-NH ₂
					PFF	GaP
					Tic	mA ₂ Bz
					tLeu	mA ₂ Bz(Dc g)
					Trp	mA ₂ Bz(Gu

n)
 Try *m*ABz
 Try(Bzl) Mapp
 Tyr Matp
 Arg(NO₂) MatpO
*p*ABz
 PaF
 PaF(Dcg)
 PaF(Mcg)
 PaF-NH₂
 PFF-NH₂
 PgF
 PzO
 Sud
 Thm
 Thm
 Tpac
 Tpac
 Tyr(Bz)O
 Me

5


[Formula IV] comprises:



10 wherein A, B, C, D, E, F, G, H, I, J, K and L are selected from the following or may be absent:

A	B	C	D	E	F	G	H	I	J	K	L
0	1	2	3	4	5	6	7	8	9	10	11
Absent or DArg	Absent or Arg	Absent or Pro	Absent or Lys	Absent or Pro	Absent or DTrp	Absent or Gln	Absent or DTrp	Absent or Phe	Absent or DTrp	Absent or Leu(r)	Absent or Leu- NH ₂ Leu
	DArg					DNMF					

5 [Formula V] comprises:




10 wherein X, A, B, C, D, E, F, G, H, I, J, and K are selected from the following or may be absent:

X	A	B	C	D	E	F	G	H	I	J	K
	-1	0	1	2	3	4	5	6	7	8	9
Absent or α -Aca 3,3Dp	Absent or Ava BAIa DmK Glt Lys Suc	Absent or DArg DNik DPaF DPzK DPzO	Absent or Arg NiK PzO	Absent or Pro	Absent or Hyp	Absent or Gly	Absent or Add Aud Ava BAIa DNMF Eac Igl Thi	Absent or DArg Ser Thr	Absent or DDab DDpr DF5F DIgl DLys DOm DPaF Nig Pac Phe	Absent or DTrp F5F Lys Nc7G Oic PaF PFF Phe	Absent or Arg Leu NiK PaF 3Pal


[Formula V] also comprises:

15





[Formula V] also comprises:

20



wherein the cyclization is via a side chain functional group other than the C-terminal residue and the residues are as described in the immediately preceding table.


25 [Formula VI] comprises the following cyclic peptides:



ACA can also be those compounds in Table 4.

35

5 [Formula VII] comprises:



wherein [ACA] is defined by Formula I or the compounds in Table 4.

10 The *in vivo* inhibitory effects of antagonists may be studied using tumor-bearing nude mice. A tumor model employing nude mice orthotopically implanted with human lung cancer cells wherein the ACA is delivered by intratracheal instillation and aerosol inhalation may be used to evaluate the efficacy and feasibility of these antagonists as a means of treating human lung cancers. Control animals without tumor implantation may
15 also be used to study the general side effects or cytotoxicity of the compounds. It is believed that aerosolized delivery or intratracheal instillation of the agents can produce effective dose accumulation in the area of lesion and reduce the overall systemic toxicity of the compounds in the animals more than when the compound is delivered by systemic administration.

20 The compounds may be administered topically, or by injection or infusion or as an oral suspension in an appropriate vehicle or as tablets, pills, capsules, caplets or the like, or preferably via intratracheal instillation or aerosol inhalation. The dosage and manner of administration will be defined by the application of the ACA and can be determined by routine methods of clinical testing to find the optimum dose. These doses are expected to
25 be in the range of 0.001 mg/Kg to 100mg/Kg of active compound.

The compounds are composed of amino acids which may form salts due to their acidic or basic nature, and any pharmacologically acceptable salt derived from the compounds described in this invention such as hydrochlorides, acetates, phosphates, maleates, citrates, benzoates, salicylates, succinates, ascorbates and the like, including HCl,
30 trifluoroacetic acid (TFA), and HOAc, are considered an extension of this invention. A common tactic in medicinal chemistry is to modify known drug substances which are peptide based to form esters or amides which exhibit greater bioavailability. Prodrugs derived from the compounds disclosed here are therefore considered an obvious extension of this invention. Methods for designing and preparing prodrugs are described in detail in
35 the medicinal chemical literature.

5 Structures and biological activities of peptides and peptide mimics related to bradykinin (BKR) are given in Table 1. Structures and biological activities of compounds not related to bradykinin (BKU) are given in Table 2. Structures and biological activities of cyclic peptides are given in Table 3. Structures of previously described known peptides which we have found to be active against cancers *in vivo* are included in Table 4. Actions
10 of selected compounds on prostate cancer cell lines are given in Table 5. Abbreviations used are as defined in Table 6.

EXAMPLES

In general, Anti-bradykinin activity was determined by the classical guinea pig
15 ileum assay and on Chinese hamster ovary (CHO) cells expressing cloned human bradykinin B2 receptors. Anti-tumor activity was determined on cultured human cancer cell lines using the standard tetrazolium (MTT) assay. No correlation between anti-bradykinin and cytolytic activity was found among the compounds, indicating that cells are not killed due to inhibition of an essential bradykinin function. Potent compounds were
20 found to stimulate apoptosis in SCLC cells, probably by abnormal activation of the intracellular MEKK pathway.

EXAMPLE I – Synthesis of peptides

Peptides were synthesized using standard solid phase synthesis methods, well
25 known in the art (Stewart and Young, *Solid Phase Peptide Synthesis*, Pierce Chemical Co., Rockford, IL, 1984) and were purified by HPLC and were characterized by amino acid analysis (AAA), thin layer chromatography (TLC) and laser desorption mass spectrometry (LDMS). Peptide amides were synthesized on methylbenzhydrylamine (MBHA) resin, which yields amides directly. Peptide methyl esters (OMe) were
30 synthesized by reaction of peptides with 2,2-dimethoxypropane (Rachele, *J. Org. Chem.* 28: 2898, 1963). Cyclic peptides were prepared on resin or in solution with PyAOP and HOAt .

5 **EXAMPLE II – Synthesis of non-peptides**

Non-peptides were synthesized by standard organic chemistry procedures well known in the art. Compounds were purified by HPLC and were characterized by analytical HPLC, TLC, and LDMS.

10 **EXAMPLE III – Synthesis of DDD and SUB dimers**

Synthesis on resin: Neutralized peptide-resin (0.05mmole) was treated with 0.15 mmole diisopropylethyl amine (DIEA) and 0.026 mmole dodecanedioyl dichloride or suberoyl dichloride in 2.5 mL dichloromethane (DCM). The suspension was mixed for 5 h, washed with DCM and ethanol and dried. The peptide dimer was cleaved from the resin
15 with HF, and the peptide was extracted and purified

Synthesis in solution: Carboxyl-derivatized amino acids or dipeptides were dissolved in dimethyl formamide (DMF) and treated with 10 equivalents of DIEA and 0.55 equivalent of dodecanedioyl dichloride or suberoyl dichloride overnight. The DMF was evaporated *in vacuo* and the resulting dimer was purified by HPLC.

20

EXAMPLE IV – Synthesis of EGS, DTP, SBEC and SUB dimers in solution

Dimerization in solution proceeded by reacting 1 equivalent of peptide monomer trifluoroacetate, an excess of DIEA and 0.55 equivalent of cross-linking reagent overnight in DMF. The cross-linking agents were purchased from Pierce (EGS dimer, ethylene glycol
25 *bis*-(succinimidylsuccinate); DTP dimer, dithio*bis* (succinimidyl propionate); SBEC dimer, *bis*[(2(succinimidooxycarbonyloxy)ethyl)sulfone; SUB dimer, disuccinimidyl suberate).

EXAMPLE V – Synthesis of Boc-N-cycloheptylglycine (Nc7G)

N-Cycloheptylglycine was synthesized by reductive amination of cycloheptanone
30 with glycine methyl ester following the procedure described in Gera *et al.*, *Immunopharmacology*. 33:174-177 (1996). The crude product was converted to the N-Boc derivative (mp, 89-90 °C).

5 **EXAMPLE VI – Synthesis of TDIM dimers**

Dimethyl tetradecyldiimidate was synthesized from tetradecanedinitrile by the method of De Abreu et al. (*Eur. J. Biochem.* 97: 379- 387, 1979. One equivalent of peptide TFA salt or other molecule having a free amino group was dissolved in DMF and stirred with 10 equivalents of DIEA and 0.7 equivalent of dimethyl tetradecyldiimidate
10 dihydrochloride overnight at room temperature. DMF was evaporated *in vacuo* and the dimer was purified. SUIM dimers were prepared similarly, using dimethyl suberimidate.

EXAMPLE VII – Synthesis of B10238: F5C-DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg (F5c-B9430)

15 B10238 was made by standard solid phase synthesis procedures, or by the acylation of B9430 with 2,3,4,5,6-pentafluorocinnamic acid in DMF, using BOP coupling in presence of excess DIEA. The product was purified by HPLC.

EXAMPLE VIII – Synthesis of M822: DDD-(DArg-F5F-Arg)₂

20 Following standard solid phase synthesis procedures, Boc-Arg(Tos) Merrifield synthesis resin was coupled with Boc-F5F, followed by coupling with Boc-DArg(Tos), using HATU as coupling agent. The peptide-resin was deprotected with TFA-DCM and neutralized with TEA. The peptide-resin was then treated with 0.55 equivalent of dodecanedioyl dichloride and 5 equivalents of DIEA in DCM overnight at room
25 temperature. After washing and drying, the resin was cleaved with anhydrous HF, using standard conditions. The peptide was extracted from the resin with 90% acetic acid and purified by preparative HPLC.

EXAMPLE IX – Synthesis of M570 Hydrochloride: F5c-OC2Y-Atmp.HCl

30 4-Amino-2,2,6,6-tetramethylpiperidine (Aldrich) was coupled with Boc-(O-2,6-dichlorobenzyl)-tyrosine, using BOP in DMF solution. The Boc protecting group was removed by TFA and the product coupled with 2,3,4,5,6-pentafluorocinnamic acid in DMF, using BOP in the presence of excess DIEA at room temperature for 3 h. The DMF was removed *in vacuo*, the product was extracted into ethyl acetate and the solvent was

5 evaporated. The residue was treated with 0.1 - 1.0 N HCl or 20% ethanolic HCl. The solvent was removed by evaporation *in vacuo* at room temperature. The residue was lyophilized from water-dioxane or crystallized from ethanol-ether.

EXAMPLE X – Synthesis of M630: Dmac-OC2Y-Matp.TFA

10 4-Methylamino-2,2,6,6-tetramethylpiperidine (Matp) was synthesized from 2,2,6,6-tetramethyl-4-piperidone (Aldrich) and methylamine by reductive amination with NaCNBH₃. The Matp was coupled with Boc-(O-2,6-dichlorobenzyl)-tyrosine, using BOP in DMF solution. The Boc protecting group was removed by TFA and the product was coupled with 4-(dimethylamino)cinnamic acid in DMF, using BOP in the presence of
15 excess DIEA at room temperature for 3 h. The DMF was removed *in vacuo*. The product was extracted into ethyl acetate and the solvent was evaporated *in vacuo*. The crude product was purified by HPLC, giving the TFA salt. The Dmac-OCTY-Matp.TFA salt can be converted to its HCL salt as in Example IX above.

20 **EXAMPLE XI – Synthesis of M638: DDD-(DArg-Igl-Arg-Matp)₂**

In sequence, Boc-Arg(Tos), Boc-Igl and Boc-DArg(Tos) were coupled to 4-methylamino-2,2,6,6-tetramethylpiperidine (Matp), using BOP as coupling agent in DMF in the presence of excess DIEA at room temperature for 3-5 h. After removal of DMF *in vacuo*, the product was extracted into ethyl acetate. After evaporation of the solvent, the
25 residue was treated with TFA-DCM to remove the Boc group. TFA was removed *in vacuo*. The DArg(Tos)-Igl-Arg(Tos)-Matp.TFA was treated with dodecanedioyl dichloride (0.55 equiv) and DIEA (5 equiv) in DCM for 5 h. The protecting groups were cleaved by HF and the lyophilized product was purified by HPLC. The M638.TFA salt was converted to its HCl salt, using 0.1 - 1.0 N HCl or 20% ethanolic HCl as in Example
30 IX above.

5 **EXAMPLE XII – Synthesis of M590: Atmp-Igl-Pac- α -Sbl-Lys-B9430**

In sequence, Boc-Igl, Boc-Pac and mono-methyl sebacate were coupled to 4-amino-2,2,6,6-tetramethylpiperidine (Atmp), using BOP coupling agent in DMF in presence of excess DIEA at room temperature for 3-5 h. DMF was removed *in vacuo* and the product was extracted into ethyl acetate. After evaporation of the solvent, the methyl ester was
10 hydrolyzed in methanol by 1N NaOH. The crude product (0.025 mmol Atmp-Igl-Pac-Sbl) was coupled to the peptide resin (0.02 mmol Lys(2-ClZ)-DArg(Tos)-Arg(Tos)-Pro-Hyp-Gly-Igl-Ser(Bzl)-DIgl-Oic-Arg(Tos)-Merrifield resin) using BOP/DIEA activation in DMF. The heterodimer peptide was cleaved from the resin with HF, using standard conditions. The peptide was extracted from the resin with acetic acid and purified by
15 preparative HPLC.

EXAMPLE XIII – Synthesis of M872: c[DArg-Arg-Eac-Ser-DF5F-Oic-Arg]

Following standard solid phase synthesis procedures, Boc-DArg(Tos) was coupled to Boc-Arg(Tos) Merrifield synthesis resin, followed in sequence by Boc-Arg(Tos), Boc-
20 Oic, Boc-DF5F, Boc-Ser(Bzl), and Boc-Eac, using HATU as coupling agent. After deprotection with TFA-DCM, the resin was cleaved with anhydrous HF using standard conditions. The peptide was extracted from the resin with 0.1% TFA-H₂O/dioxane and lyophilized. The peptide trifluoroacetate was cyclized with three equivalents of PyAOP and HOAt and 20 equivalents of DIEA in DMF at a concentration of 10⁻³ M. After
25 removal of the solvent under reduced pressure, the product was lyophilized from dioxane-H₂O and purified by HPLC.

EXAMPLE XIV – Synthesis of M678: (Dns-DArg-Igl-Arg)₂-DDA

In sequence, Boc-Arg(Tos), Boc-Igl and Boc-DArg(Tos) (2 equivalents) were
30 coupled to 1,10-decanediamine using BOP as a coupling agent in DMF in presence of excess DIEA at room temperature for 3-5 h. DMF was removed *in vacuo* and the product was extracted into ethyl acetate. The solvent was evaporated *in vacuo* and the residue was treated with TFA/DCM to remove the Boc group. TFA was removed *in vacuo*, and the

- 5 product was treated with dansyl chloride (2 equivalents) and an excess of DIEA in DCM for 5 h. The Tos groups were cleaved by HF and the crude product was purified by HPLC.

EXAMPLE XV – Synthesis of M290: BTAC-(2-Nal-Atmp)₃

- 10 The benzene-1,3,5-*tris*-carbamido- ϵ -caproic acid linker was made from 1,3,5-benzenetricarboxylic acid and N-Boc- ϵ -caproic acid methyl ester, using the BOP coupling method. The methyl ester was hydrolyzed in methanol by 1N NaOH. The product (1 equivalent BTAC) was coupled to 2-Nal-Atmp (3 equivalents) in DMF, using HATU as coupling agent. The solvent was removed *in vacuo*, and the residue was purified by HPLC.
- 15 The BTAC-(2-Nal-Atmp)₂-OH was also isolated as a by-product.

EXAMPLE XVI – Synthesis of M1040: EDTA-(OC2Y-ATMP)₄

- Boc-(O-2,6-dichlorobenzyl)-tyrosine was coupled with 4-amino-2,2,6,6-tetramethylpiperidine overnight in DMF, using BOP as coupling agent in the presence of
- 20 DIEA. After removal of DMF *in vacuo*, the residue was extracted into ethyl acetate and treated with TFA/DCM to cleave the Boc group. The TFA/DCM was evaporated *in vacuo* and the product (OCTY-ATMP) was lyophilized from dioxane/water.
- Ethylenediaminetetraacetic acid (0.25 equivalent EDTA) was coupled with OC2Y-ATMP trifluoroacetate (1 equivalent) in DMF, using BOP as coupling agent in the presence of
- 25 DIEA. The solvent was removed *in vacuo* and the residue was purified by HPLC.

EXAMPLE XVII - Assay of anti-bradykinin activity on guinea pig ileum

- Male Hartley guinea pigs that had been deprived of food overnight were sacrificed, and sections of terminal ileum, 25 mm in length, were dissected, attached to tissue holders
- 30 and immersed in 10 ml tissue baths containing Krebs' solution bubbled with 95%O₂/5%CO₂. Tissues were placed under 1 g tension and incubated for 1 h equilibration. Concentration-effect curves were constructed to bradykinin in the absence and presence of new compounds. Bradykinin showed pD₂ = 7.4, and antagonist B9430 showed pA₂ = 7.9.

5

EXAMPLE XVIII - Assay of anti-bradykinin activity on cloned human B2 receptors

Chinese hamster ovary cells containing cloned and expressed human bradykinin B2 receptors were grown in cell cups of the Cytosensor microphysiometer in Ham's F-12 medium supplemented with sodium pyruvate and 10% FBS (Gibco 11765-054). For assay
10 the cells were transferred to Ham's F-12 without bicarbonate or serum (Gibco 21700-075) and placed in the Cytosensor. Concentration-response curves were constructed to bradykinin in the presence or absence of new compounds. Bradykinin showed $pD_2 = 11$, and antagonist B9430 showed $pA_2 = 10.5$

15 **EXAMPLE XIX - Colorimetric tetrazolium assay for cell survival**

Cell growth and survival were measured by a rapid colorimetric assay based on the tetrazolium salt MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide) (Mosmann, *J. Immunol. Methods* 65: 55-63, 1983, with minor modifications). Briefly, 1,000 normal lung fibroblasts or normal epithelial BEAS-2B cells, 1,000 or 5,000 viable
20 non-SCLC cells or 10,000 viable SCLC cells were plated in 100 μ L of growth medium in 96-well flat-bottomed microtiter plates. Cells were incubated overnight to allow recovery. Compounds to be tested were added to the cells in triplicate in a range of concentrations and the cells were incubated at 37 $^\circ$ C, 5% CO₂, with 100% humidity. Control cells were treated in the same way without antagonists. All wells had a final volume of 200 μ L. Plates
25 were incubated for 4 days, allowing sufficient time for cell replication and compound-induced cell death to occur. On day 5, 25 μ L of a 2 mg/mL solution of MTT (Sigma) dissolved in RMPI-1640 was added to each well. The plate was incubated for 4 h at 37 $^\circ$ C. The supernate was removed and the blue formazan complex was dissolved by adding 100 μ L of 0.02 N HCl in 75% isopropanol to all wells. Absorbance was immediately
30 determined using a scanning multiwell plate reader. B9870 caused 50% cell death at a concentration of 0.15 μ M under these conditions.

5 **EXAMPLE XX - Measurement of apoptosis in cultured cells**

Apoptosis, also known as programmed cell death, is the phenomenon by which a cell dies following a series of gene-mediated events, in response to a wide range of intracellular and extracellular agents. Apoptosis, a counterpart of mitosis, plays an important role in the development and homeostasis of many organisms and tissues. It serves to regulate cell numbers, to shape developing organisms and as a defense against potentially harmful agents. Apoptosis is not the only mode of cell death. Necrosis is a type of cell death which is nonspecific and frequently occurs when cells are exposed to high doses of toxic agents. Such exposure usually results in the loss of ionic homeostasis. Unlike apoptosis, necrosis does not seem to be genetically influenced.

15 Apoptotic and necrotic cells have different appearances which allow them to be distinguished microscopically. Necrotic cells and their mitochondria swell, the cell membrane eventually ruptures, and internal organelles become distended. As a result of the membrane rupture, inflammation occurs in the surrounding tissue. In contrast, the nuclei of apoptotic cells become fragmented into several smaller nuclear bodies, which are quickly recognized by phagocytes and engulfed, and no inflammatory response occurs. Therefore, it is useful to develop chemotherapeutics which induce apoptosis, rather than necrosis, in order to avoid inflammation and the toxic agents which are often released from necrotic tumor cells.

We have used differential fluorescent dye uptake and cellular morphology to distinguish viable and dead cells with apoptotic and/or necrotic morphologies. We have used Rhodamine 123 to stain active mitochondria in viable cells, Hoechst 33324 to stain DNA in both viable and dead cells, and Propidium Iodide to stain DNA in dead cells. These cell subpopulations may be distinguished by the different manners in which they take up the fluorescent probes. The dead apoptotic and necrotic subpopulation, which has lost its membrane potential and organelle function, takes up Propidium Iodide and Hoechst 33324. Since the cells in this subpopulation are dead, the mitochondria are not active and thus there is little or no uptake of Rhodamine 123. Under the fluorescence microscope with a DAPI filter, nuclei in these cells appear pinkish in color due to the mixing of both Propidium Iodide and Hoechst 33324 dyes. Necrotic cells have intact nuclei while

5 apoptotic cells have fragmented multi-nucleated bodies. In contrast, the viable apoptotic subpopulation has an intact membrane but inactive mitochondria. As a result, the fragmented multi-nucleated bodies (a hallmark of apoptotic cells) in these cells take up only Hoechst 33324, which gives them a blue appearance under the fluorescence microscope, but are unable to take up Propidium Iodide or Rhodamine 123. The
10 subpopulation of viable cells has both intact cell membranes and active mitochondria. These cells take up both Hoechst 3324 and Rhodamine 123. Microscopically these cells appear to have single normal blue nuclei when examined with a DAPI filter and bright green mitochondria when examined with a FITC filter.

15 **EXAMPLE XXI – Inhibition of tumor growth *in vivo* in nude mice**

Representative peptide and non-peptide compounds having high *in vitro* cytotoxic activity were tested against implanted tumors *in vivo*. Athymic nude mice were implanted subcutaneously with either single cell suspensions (2 million SCLC cells or 1 million NSCLC cells) or with small fragments (3 x 3 mm) of tumors minced from previously
20 grown nude mouse heterotransplants. On the seventh day after tumor implantation groups of 5 mice bearing implants were injected intraperitoneally daily with the compounds being tested at 1, 5, or 10 mg/kg/day; control animals were injected with an equal volume of isotonic saline. Tumor size was measured with a caliper three times per week. Tumor volume was calculated by the formula:

25
$$\text{Volume (cc)} = \pi \times (\text{length}) \times (\text{width})^2 / 6$$

Results of representative *in vivo* tests are given in Figs. 1-8. For comparison, bradykinin antagonist peptide dimers B9870 and B10054 caused marked inhibition of growth of the SCLC line SHP-77 at a dose of 5 mg/kg/day.

30 **EXAMPLE XXII - Data**

Examples of peptides and peptide mimics related to the C-terminal part of bradykinin antagonist peptides and their biological activities on cancer cells and bradykinin responses are given in Table 1.

5 Many compounds not directly related to the structure of bradykinin were synthesized and tested for anti-tumor and anti-bradykinin activity. These are listed in Table 2.

Cyclic peptides related to bradykinin and bradykinin mimics are reported in Table 3, along with their biological activity on cancer cells and anti-bradykinin activity.

10 Structures of previously described known peptides which have been found to be active against cancers *in vivo* are included in Table 4.

Cytotoxic activity *in vitro* of compounds M570 and M590 against various standard strains of prostate cancer is reported in Table 5.

15 Standard abbreviations were used for natural amino acids. For non-natural amino acids, derivatizing groups and other chemicals, the abbreviations listed in Table 6 are used.

Table 1. ACTIVITIES OF PEPTIDES RELATED TO BRADYKININ STRUCTURE

NUMBER	STRUCTURE	MTT ^a	GPI ^b	HUMAN ^c
BK ^d	Arg-Pro-Pro-Gly-Phe-Ser-Pro-Phe-Arg			
B9430 ^d	DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg	120	8.2	
B9870-2 ^d	SUIM-(DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg)	0.15	8.4	
B8838	DArg-Arg-Pro-Hyp-Gly-CpG-Ser-DCpG-CpG-Arg	--	7.0	
B8840	DArg-Arg-Pro-Hyp-Gly-Phe-Ser-DCpG-CpG-Arg	--	6.8	
B8858	DArg-Arg-Pro-Hyp-Gly-Thi-Ser-CpG-DCpG-DArg	--	5.2	
B8994	DArg-Arg-Pro-MeP-Gly-CpG-Ser-DCpG-CpG-Arg	--	--	
B9074	Dhq-DArg-Arg-Pro-Hyp-Gly-CpG-Ser-DCpG-CpG-Arg	--	6.3	
B9126	Aaa-DArg-Arg-Pro-Hyp-Gly-(D,L)DMF-Ser-DTic-Oic-Arg	--	6.4	
B9126-2	Aaa-DArg-Arg-Pro-Hyp-Gly-(D,L)DMF-Ser-DTic-Oic-Arg	--	7.3	
B9224-2	Aca-DArg-Arg-Pro-Hyp-Gly-Thi-Ser-(D,L)Igl-Oic-Arg	--	8.4	
B9882	α -Sub-Lys(ϵ Flu)-DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg	--	--	
	^L DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg	--	--	
B9914	Oic-Arg	--	Wk	
B9916	DIgl-Oic-Arg	--	Wk	
B9490	Dcg-Digl-Oic-Arg	>60	Wk	
B9918	Ser-Digl-Oic-Arg	--	--	
B9920	Igl-Ser-DIgl-Oic-Arg	--	--	

B9922	Gly-Igl-Ser-DIgl-Oic-Arg	--	Wk	
B9924	Hyp-Gly-Igl-Ser-Digl-Oic-Arg	--	--	
B9926	Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg	--	--	
B9950	α -Lys-DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg	8	--	
	Sub-Arg-DNMF-DTrp-Phe-DTrp-Leu			
B9956	α -DDD-(Lys-DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg) ₂	--	--	
B9960	DArg-Arg-Nig-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg	--	7.7	
B9966	DArg-Arg-NMF-Hyp-Gly-Thi-Ser-DIgl-Oic-Arg	--	6.9	
B10010	DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Nc7G-Arg	--	7.7	
B10014	DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Nc6G-Arg	--	7.6	
B10054	DDD-(Lys-DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg) ₂	0.3	7.1	
B10062	DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg-NH ₂	Inact	7.1	
B10082	SUIM-(DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg-NH ₂) ₂	0.7	7.2	
B10084	BAPG-DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg	>20	8.1	
B10088	DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg-Eac-Eac-DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg	4	7.1	
B10092	(Gun) ₂ -BAPG-DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg	>20	8.7	
B10098	(DArg-Arg-Pro-Hyp) ₂ -Dpr-Igl-Ser-DIgl-Oic-Arg	20	5.3	
B10100-2	TDIM-(DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg) ₂	1	8.0	
B10100-1	Moti-DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg	4	7.8	
B10104-2	TDIM-(DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DIgl-Oic-Arg) ₂	4	8.0	
B10104-3	Moti-DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DIgl-Oic-Arg	20	8.1	
B10160	Leu-DTrp-Phe-DTrp-DNMF-Eac ₂ -DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg	10	6.3	
B10162	Leu-Leu-DTrp-Phe-DTrp-DNMF-Eac ₂ -DArg-Arg-Pro-Hyp-Gly-Igl-Ser-Digl-Oic-Arg	7	6.3	
B10198-1	DDD-(Eac-Arg-Digl-Oic-Arg) ₂	--	5.7	
B10198-2	DDD-(Eac-Arg-Digl-Oic-Arg) ₂	15	--	
B10200	DDD-(Eac-Eac-Arg-Digl-Oic-Arg) ₂	16	5.8	
B10238	F5c-DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg	150	8.1	
B10252	EGS-(DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg) ₂	25	8.3	
B10282	Arg-Pro-Pro-Gly-Phe-Thr-DTic-Oic-Arg	--	7.3	
B10284	Arg-Pro-Pro-Gly-Phe-Thr-DTic-Oic-NH ₂	--	7.7	
B10382	DArg-PzO-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg	--	--	
B10384	DNiK-PzO-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg	--	--	
B10386	DDD-(DmK-PzO-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg) ₂	--	--	
B10388	DNiK-PzO-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg	--	--	
B10390	DNiK-PzO-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg	--	--	
B10392	DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-PFF-Arg	--	--	
B10394	F5c-DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-PFF-Arg	--	--	
B10396	F5c-DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-PFF-Arg	--	--	
M2	Dcg-D-2-Nal-Arg	--	--	
M10	Dcg-2Nal-Arg	--	4.8	--
M20	Gun-2-Nal-Arg	--	4.6	
M42	Gun-Eac-DIgl-Oic-Arg	--	5.0	
M68	Dcg-BtA-Arg	--	4.9	
M70	Dcg-Igl-Arg	--	4.8	

M78	Dcg-Apa-Arg	--	5.6	
M84	Dcg-Ile-Arg	--	5.1	
M86	Dcg-Ac6c-Arg	--	5.0	
M88	Gun-Ica-Arg	--	4.7	
M94	Dcg-Aic-Arg	--	Wk	
M96	Dcg-(D,L)Atc-Arg	--	4.7	
M118-1	Ac-PaF(Mcg)-Arg	--	4.9	
M118-3	Ac-PdF-Arg	--	5.4	
M124	Dcg-Gly-Cmp-Arg	--	4.7	
M128	Dcg-Gly-Oic-Arg	--	Wk	
M130	Dcg-F5F-Arg	20	4.8	
M132	F5bz-F5F-Arg	60	Wk	
M134	Dcg-Trx-Arg	--	4.9	
M142	Ac-PaF(Sin)-Arg	--	Wk	
M146-1	Ac-PaF(Mcg)- <i>p</i> -ABz-Arg	--	Inact	
M146-2	Ac-PaF(Dcg)- <i>p</i> -ABz-Arg	--	4.9	
M148	F5c- <i>p</i> -ABz-Arg	--	5.1	
M160	Ste-2-Nal-Arg	--	Wk	
M176	F5c- <i>p</i> ABz-2Nal-Arg	Inact	5.4	--
M196	F5c-Gly- <i>m</i> ABz-2Nal-Arg	Inact	5.1	--
M198	Ac-Pac-Gly- <i>m</i> -Abz-2-Nal-Arg	--	5.1	
M200-1	Mcg-Pac-Gly- <i>m</i> -ABz-2-Nal-Arg	--	Inact	
M200-2	Dcg-Pac-Gly- <i>m</i> -ABz-2-Nal-Arg	--	4.9	
M216	F5c- <i>p</i> -APa-Arg	>180	--	
M226	DDD-(Arg-Dlgl-Oic-Arg) ₂	35	5.7	Inact
M232-1	Dcg-Atpc-Arg	--	4.7	
M232-3	Dcg-2-Nal-Atpc-Arg	--	5.1	
M346	Dcg- <i>p</i> -Amb-Arg	--	4.6	
M348	F5c- <i>p</i> -Amb-Arg	--	4.7	
M352	F5c- <i>p</i> -Amb-APa-Arg	--	4.7	
M370	F5c-Arg	--	4.8	
M372	F5c-APb-Arg	--	4.6	
M374	Tfmc-Arg	--	4.6	
M380	F5c-Tyr-Arg	--	Inact	
M382	F5c-Tic-Arg	--	4.7	
M388	F5c-Lys{(CH ₃) ₃ }-Arg	--	4.9	
M392	F5c-Ana-Arg	--	4.5	
M394	F5c-Bip-Arg	--	4.7	
M398	F5c-Pac-Arg	--	Inact	
M400	DDD-(<i>p</i> ABz-2Nal-Arg) ₂	22	5.1	11.5
M406	Arg-Eac-Dlgl-Ana-Arg	--	Inact	
M410	F5c-Phe-Arg	--	5.4	
M412	F5c- <i>m</i> -APa-Arg	--	5.8	
M416	F5c-3-Pal-Arg	--	Wk	
M420	F5c-hPhe-Arg	60	7.0	10.9
M424	F5c-Thi-Arg	--	4.6	
M426	F5c-Trp-Arg	--	Inact	
M442	F5c-Oic-Arg	--	--	
M446	F5c-2Nal-Arg	60	4.7	9
M450	F5c-2Nal-Arg-NH ₂	26	4.9	Inact
M484	DDD-(Pac-2Nal-Arg) ₂	25	Inact	Inact
M494	DDD-(Lys-Pac-Gly- <i>m</i> ABz-2Nal-NH ₂) ₂	33	5.1	Inact
M498	DDD-(Pac-2Nal-Arg-NH ₂) ₂	24	4.9	Inact
M500	DDD-(<i>p</i> ABz-2Nal-Arg-NH ₂) ₂	40	0	11.4

M504	DDD-(Pac-2Nal-DArg-NH ₂) ₂	11	5.4	Wk
M508	DDD-(DArg-2Nal-Arg) ₂	23	Inact	--
M510	DDD-(DArg-2Nal-Arg-NH ₂) ₂	8	Inact	11
M512	F5c-OC2Y-Arg	70	5.7	11
M516	DDD-(DArg-Arg-Aud-Pac-2Nal-Arg) ₂	1.4	0	Ag
M518	DDD-(DArg-OC2Y-Arg) ₂	15	Wk	10
M520	F5c-OBS-Arg	Inact	6.1	7
M528	F5c-MBC-Arg	Inact	Inact	
M540	Pya-hPhe-Arg	>100	Wk	
M542	Dca-hPhe-Arg	80	Wk	
M550	F5c-OBT-Arg	80	Inact	Ag
M552	DDD-(p-ABz-hPhe-Arg) ₂	Inact	Inact	
M554	DDD-(DArg-hPhe-Arg) ₂	--	5.1	
M558	Dcg-hPhe-Arg	100	Wk	
M560	DDD-(DArg-hPhe-Arg-NH ₂) ₂	50	Wk	
M564	DDD-(DArg-OBS-Arg) ₂	--	Wk	
M590	Atmp-Igl-Pac- α -Sbl-Lys-B9430	4.5	7.5	Inact
M598	DDD-(Arg-DIgl-Oic-Arg-OMe) ₂	13	--	10
M600	α -DDD-(Lys-B9430-OMe) ₂	1.2	6.4	Ag
M608	DDD-(Eac-Arg-DIgl-Oic-Arg-OMe) ₂	13	--	Inact
M612	F5c-DArg-hPhe-Arg	Inact	Wk	
M676	DDD-(DArg-Arg-Eac-Ser-DF5F-Nc7G-Arg) ₂	--	--	
M682	F5c-Lys(F5bz)-Arg	--	4.9	
M686	F5c-NMF-Arg	29	5.6	
M688	F5c-Dpr(Fbz)-Arg	--	Inact	
M690	F5c-Dpr(Paa)-Arg	--	4.8	
M692	DDD-(DArg-Arg-Aud-Pac-hPhe-Arg) ₂	--	Wk	
M696	F5c-DArg-Eac-2Nal-Arg	Inact	5.1	--
M698	F5c-DArg-Arg-Aud-Pac-2Nal-Arg	7.1	Wk	--
M706	Cin-hPhe-Arg	Inact	Wk	
M708	Ppa-hPhe-Arg	Inact	5.8	
M710	DDD-(DArg-Arg-Aud-Pac-2Nal-DArg-NH ₂) ₂	1.7	Wk	
M714	F5c-PCF-Arg	Inact	Wk	
M718	F5c-PFF-Arg	9	5.1	
M720	F5c-PaF(Ppa)-Arg	Inact	4.8	
M726	D-Arg-Arg-Aud-PaF(F5c)-Arg	Inact	5.6	
M728	DDD-(DArg-Arg-Aud-PaF(F5c)-Arg) ₂	4	5.3	
M730	F5c-DhPhe-Arg	--	4.7	
M732	F5c-PNF-Arg	--	4.9	
M734	DDD-(DArg-Arg-Aud-Pac-PaF(Fbz)-Arg) ₂	1.8	5.3	
M738	F5c-DArg-Eac-hPhe-Arg	Inact	5.3	
M746	DDD-(Pac-hPhe-Arg) ₂	Inact	Inact	
M752-2	Pac-hPhe-Arg	Inact	Wk	
M752-5	Aaa-Ser-Pac-hPhe-Arg	Inact	Wk	
M752-6	Aaa-Pac-hPhe-Arg	Inact	Wk	
M754	Aaa-DPhe-hPhe-Arg	Inact	4.6	
M756	DDD-(DPhe-hPhe-Arg) ₂	18	5.2	
M758	Saa-hPhe-Arg	--	Wk	
M764	Aaa-DTic-hPhe-Arg	--	4.8	
M766	F5c-DArg-Arg-Aud-DTic-hPhe-Arg	--	5.1	
M770	DDD-(DArg-Arg-Aud-DTic-hPhe-Arg) ₂	8	Inact	
M772	Aaa-DIgl-hPhe-Arg	--	4.9	
M774	F5c-DArg-Arg-Aud-DIgl-hPhe-Arg	8	Inact	
M776	DDD-(DIgl-hPhe-Arg) ₂	30	5	
M778-1	Pcc-hPhe-Arg	--	Wk	
M780	Mca-hPhe-Arg	--	Wk	

M782	Cca-hPhe-Arg	--	Wk
M784	Ac-OC2Y-Arg	Inact	Wk
M786	DDD-(DArg-Arg-Aud-DIgl-hPhe-Arg) ₂	3.2	Wk
M788	F5c-DArg-Arg-Aud-DTic-Oic-Arg	9	5
M790	DDD-(DArg-Arg-Aud-DTic-Oic-Arg) ₂	1.7	Inact
M792	F5c-DArg-Arg-Eac-Ser-DTic-Oic-Arg	>100	4.9
M794	DDD-(DArg-Arg-Eac-Ser-DTic-Oic-Arg) ₂	21	Inact
M796	F5c-DArg-Arg-Eac-Ser-DF5F-Oic-Arg	31	6.5
M802	F5c-Lys-Ser-DF5F-Oic-Arg	Inact	6.3
M804	DDD-(DArg-Arg-Eac-Ser-DF5F-Oic-Arg) ₂	7.3	7.7
M806	Ava-Igl-Ser-DF5F-Oic-Arg	Inact	5.6
M808	DDD-(Lys-Ser-DF5F-Oic-Arg) ₂	30	6.9
M810	F5c-F5F-Arg	40	4.6
M812	F5c-PFF-Arg-NH ₂	15	Wk
M814	Ppa-PFF-Arg	Inact	Wk
M816	Dpa-PFF-Arg	52	4.6
M818	DDD-(DArg-PFF-Arg-NH ₂) ₂	60	Wk
M820	DDD-(DArg-PFF-Arg) ₂	43	Inact
M822	DDD-(DArg-F5F-Arg) ₂	25	Mixed
M826	F5c-MFF-Arg	76	Inact
M828	F5c-3,4F2F-Arg	--	Wk
M838	F5c-DArg-Arg-Aud-DIgl-PFF-Arg	7.4	5.1
M842	DDD-(DArg-Arg-Aud-DIgl-PFF-Arg) ₂	1.4	Inact
M844	DArg-Arg-Aud-DIgl-PFF-Arg	12	Wk
M846	DDD-(DArg-Arg-Aud-DF5F-Oic-Arg) ₂	2	7.1
M852	F5c-DArg-Arg-Eac-Ser-DIgl-Oic-Arg	Inact	5.9
M854	DDD-(DArg-Arg-Eac-Ser-DIgl-Oic-Arg) ₂	7.3	5.9
M856	F5c-DArg-Arg-Aud-Ser-DIgl-Oic-Arg	21	5.4
M858	DDD-(DArg-Arg-Aud-Ser-DIgl-Oic-Arg) ₂	4	6.3
M860	F5c-DArg-Arg-Add-Ser-DIgl-Oic-Arg	6	5.4
M862	DDD-(DArg-Arg-Add-Ser-DIgl-Oic-Arg) ₂	1.3	5.6
M864	DDD-(DArg-Arg-Add-Ser-DIgl-PFF-Arg) ₂	1.8	Inact
M868	Ac-Darg-Arg-Aud-DF5F-Oic-Arg	55	6.5
M888	F5c-DArg-Arg-Aud-Ser-DF5F-Oic-Arg	12.5	6.6
M890	DDD-(DArg-Arg-Aud-Ser-DF5F-Oic-Arg) ₂	1.7	5.5
M922	DDD-(DNiK-Arg-Eac-Ser-DF5F-Oic-Arg) ₂	--	--
M926	ζ-SUB-(ApC-F5F-Arg) ₂	Inact	--
M930	α-DDD-(ApC-F5F-Arg) ₂	Inact	--
M932	DDD-(DArg-Arg-Eac-Ser-DIgl-PFF-Arg) ₂	6.0	--
M936	DDD-(DNiK-PzO-Eac-Ser-DF5F-Oic-Arg) ₂	--	--
M944	DDD-(DArg-Arg-Eac-Ser-DF5F-PFF-Arg) ₂	6.7	--
M946	F5c-DArg-Arg-Eac-Ser-DF5F-PFF-Arg	--	--
M950	α-DDD-(K-DArg-Arg-Eac-Ser-DF5F-Oic-Arg) ₂	6.7	--
M952	DDD-(DmK-DArg-Arg-Eac-Ser-DF5F-Oic-Arg) ₂	--	--
M954	Aaa-DArg-Arg-Eac-Ser-DF5F-Oic-Arg	10	--
M956	Aaa-DArg-Arg-Aud-Ser-DF5F-Oic-Arg	14	--
M958	F5bz-DArg-Arg-Aud-Ser-DF5F-Oic-Arg	18	--
M960	Aca-DArg-Arg-Aud-Ser-DF5F-Oic-Arg	21	--
M964	33Dp-DArg-Arg-Aud-Ser-DF5F-Oic-Arg	4-8	--
M968	Dmac-DArg-Arg-Aud-Ser-DF5F-Oic-Arg	15	--
M972	F5pa-DArg-Arg-Aud-Ser-DF5F-Oic-Arg	--	--
M974	DDD-(PzO-F5F-Arg) ₂	--	--
M976	DDD-(DNiK-F5F-Arg) ₂	--	--
M978	DDD-(DPzK-F5F-Arg) ₂	--	--

M980	DDD-(DPzO-F5F-Arg) ₂	--	--
M1024	SUB-(DArg-Arg-Eac-Ser-DF5f-Nc7G-Arg) ₂	14	7.2
M1026	DTP-(DArg-Arg-Eac-Ser-DF5F-Nc7G-Arg) ₂	70	6.9
M1028	SBEC-(DArg-Arg-Eac-Ser-DF5F-Nc7G-Arg) ₂	28	6.7
M1030	EGS-(DArg-Arg-Eac-Ser-DF5F-Nc7G-Arg) ₂	51	7.0
M1034	DDD-(DArg-F5F-DArg-NH ₂) ₂	--	--
M1036	DDD-(DArg-F5F-DArg) ₂	40	5.4
M1038	ε-SUB-(Lys-DArg-Arg-Eac-Ser-DF5F-Nc7G-Arg) ₂	--	6.4
M1042	Aca-DArg-Arg-Eac-Ser-DF5F-Oic-Arg	--	--
M1044	Gun ₂ -BApg-DArg-Arg-Eac-Ser-DF5F-Oic-Arg	--	--
M1046	(F5c-DArg-Igl-Arg) ₂ -DDA	--	--

5

Footnotes:

^a ED₅₀ for killing of SCLC strain SHP-77 *in vitro*, μM.

10 ^b pA₂ for bradykinin antagonist activity on isolated guinea pig ileum. The pD₂ of bradykinin is 7.4 on ileum. Higher numbers indicate higher potency.

^cpA₂ for bradykinin antagonist potency on cloned human B2 receptors, pM. The pD₂ for bradykinin is 11. Higher numbers indicate higher potency.

^d Data included for comparison

15 Inact = inactive; Mixed = showing both agonist and antagonist activity; Wk = weak

Table 2. ACTIVITIES OF COMPOUNDS NOT RELATED TO BRADYKININ

NUMBER	STRUCTURE	MTT ^a	GPI ^b	HUMAN ^c
B9948	Arg-DNMF-DTrp-Phe-DTrp-Leu	2.8	Wk	
B10222	DNMF-DTrp-Phe-DTrp-LeuΨ(CH ₂ NH)Leu-NH ₂	6.0	5.2	
B10224-1	α-DDD-(Lys-DNMF-DTrp-Phe-DTrp-LeuΨ(CH ₂ NH)Leu-NH ₂) ₂	13	--	
B10224-2	α-DDD-(Lys-DNMF-DTrp-Phe-DTrp-LeuΨ(CH ₂ NH)Leu-NH ₂) ₂	7	--	
B10228	DDD-(DNMF-DTrp-Phe-DTrp-LeuΨ(CH ₂ NH)Leu-NH ₂) ₂	40	Wk	
B10242	Arg-Pro-Lys-Pro-DTrp-Gln-DTrp-Phe-DTrp-LeuΨ(CH ₂ NH)Leu-NH ₂	40	5.6	
B10244	DArg-Arg-Pro-Lys-Pro-DTrp-Gln-DTrp-Phe-DTrp-LeuΨ(CH ₂ NH)Leu-NH ₂	12	5.4	
B10246	DArg-Pro-Lys-Pro-DTrp-Gln-DTrp-Phe-DTrp-LeuΨ(CH ₂ NH)Leu-NH ₂	12	6.1	
>278	F5c-Iqa-Atmp	9	5.3	--

M8	Gun-Eac-D2Nal-PgF	--	Inact	
M12	Dcg-Igl-Aqu	20	5.0	--
M18	Dcg-2Nal-Aqu	30	6.6	--
M26	Gun-2Nal-GaP	--	4.9	
M30	Dcg-2Nal-Apa	--	5.4	
M32	Gun-2Nal-Apa	--	4.8	
M36	Dcg-D2Nal-Apa	--	5.0	
M38	Gun-D2Nal-Apa	--	4.9	
M62	Dcg-2Nal-Ama	--	4.8	
M64	Dcg-2Nal-APa-Sud	--	Ag	
M72-1	Dcg-Igl-Apa	--	4.7	
M72-2	Dcg-Igl-APa(anisyl)	--	4.6	
M76	Dcg-2Nal- <i>m</i> ABz	--	4.9	
M92-1	Dcg-2Nal- <i>m</i> A ₂ Bz	--	5.0	
M92-2	Dcg-2Nal- <i>m</i> A ₂ Bz(Gun)	--	4.8	
M92-4	Dcg-2Nal- <i>m</i> A ₂ Bz(Dcg)	--	5.0	
M104	Dcg-2Nal-3Pal	--	4.9	
M112	Dcg-D2Nal- <i>m</i> ABz	--	5.1	
M120	Dcg-2Nal- <i>p</i> ABz	--	4.7	
M122-1	Mcg-APa- <i>m</i> ABz	--	5.0	
M122-2	Dcg-APa- <i>m</i> ABz	--	4.6	
M136	Sin-F5F-3Pal	--	Inact	
M162	Dcg-2Nal-Asp	--	4.9	
M168-1	2Nap-PaF(Mcg)	--	4.8	
M168-2	2Nap-PaF(Dcg)	--	4.7	
M172	Inp-Dpr(Dcg-2Nal)	--	4.9	
M174	Dcg-Asp-Aqu	--	Inact	
M180	F5c- <i>p</i> ABz-2Nal	--	5.1	
M188B	Dcg-2Nal-Asp(Aqu)	--	5.5	
M202	F5c-Gly- <i>m</i> ABz-2Nal	--	5.1	
M204	Ac-Pac-Gly- <i>m</i> ABz-Nal	--	5.0	
M218	2Nal-Atmp	Inact	4.8	
M222	Dcg-2Nal-Atmp	15	6.8	Inact
M228-2	Dcg(Me)-2Nal-Atmp(Me)	15	7.6	--
M236	Dcg-Igl-Atmp	>50	4.7	--
M240	Dcg-F5f-Atmp	32	5.1	--
M244A	Dcg-2Nal-Atpm	12	5.0	
M244B	Dcg-2Nal-Atpc	Inact	4.9	
M246	Dcg-D2Nal-Atmp	>50	5.7	
M248	F5c-2Nal-Atmp	3.2	6.2	--
M250	Aca-2Nal-Atmp	--	5.2	
M252	Dhq-2Nal-Atmp	Inact	4.8	
M254	TDIM-(2Nal-Atmp) ₂	5	5.1	--
M254-1	TDIM-(2Nal-Atmp) ₂	5	5.1	
M254-2	TDIM-(2Nal-Atmp) ₂	5	5.8	
M258	Dcg-Igl-Atp	--	5.0	
M262	Dcg-D2Nal-Atmp	6	5.3	
M264	Dcg-Trp-Atmp	16	4.7	--
M266	Dcg-APa-Atmp	42	4.8	--
M268	F5c-2Nal-Tpac	10	5.2	--
M270	Dcg-2Nal-Tpac	--	6.1	
M272	Dpa-2Nal-Atmp	9	5.2	--

M274	Sin-2Nal-Atmp	36	4.7	--
M276	Dca-2Nal-Atmp	4.6	5.3	--
M280	TDIM-(Igl-Atmp) ₂	6	5.2	--
M280-1	Ctim-Igl-Atmp	21	Wk	
M280-2	TDIM-(Igl-Atmp) ₂	6	5.2	
M286	Dtp-(2Nal-Atmp) ₂	24	5.1	--
M288	Boc-2Nal-Atmp	Inact	5.2	--
M288A	Boc-2Nal-Atmp	>85	5.2	
M290-1	Btac-(2Nal-Atmp) ₂	>60	Wk	
M290-2	Btac-(2Nal-Atmp) ₃	20	Wk	
M292	Pac-Igl-Atmp	40	Wk	
M294	DDD-(Pac-Igl-Atmp) ₂	1.8	Inact	10.3
M296	Pya-Bip-Atmp	15	Wk	
M302	Atep-2Nal-Atmp	3.5	5.2	
M304	TDIM-(2Nal-Dmm) ₂	4.2	5.9	--
M306	Gbz-2Nal-Atmp	>100	--	
M308	Pac-2Nal-Atmp	>75	5.0	
M310	DDD-(Pac-2Nal-Atmp) ₂	1.2	5.1	Inact
M312	Tfmc-2Nal-Atmp	3.2	5.5	--
M314	F5c-2Nal-Aqd	25	4.6	
M316	F5c-Tyr-Atmp	50	--	
M318	F5c-Tyr(Bzl)-Atmp	3.6	5.1	--
M320	F5c-Oic-Atmp	13	--	--
M322	F5c-Tic-Atmp	7.6	--	--
M324	Dmac-2Nal-Atmp	3	5.2	--
M336-1	Dcg-2Nal-Asp-(R,S)Aqu	--	5.1	
M336-2	Dcg-2Nal-Asp-(R,S)Aqu	--	5.4	
M340	Dcg-Pac-Gly- <i>m</i> ABz-2Nal	--	4.8	
M342	Dcg-2Nal-Asp-Atmp	--	5.4	
M350	Dcg-2Nal-Glu-Atmp	--	5.0	
M354	Dcg-2Nal-PgF	--	5.2	
M362	Dcg- <i>p</i> APa-Asp-Atmp	--	4.9	
M364	F5c- <i>p</i> APa-Asp-Atmp	--	4.7	
M368	Tfmc- <i>p</i> APa-Asp-Atmp	--	4.7	
M396	F5c-2Nal-Cys(SO ₃ H)-Atmp	--	5.0	
M408	Pya-2Nal-Cyh	22	4.6	11.5
M418	F5c-BtA-Atmp	3.8	7.0	10.3
M422	Pya- <i>p</i> ABz-2Nal	52	--	Inact
M428	Pya-Gly- <i>m</i> ABz-Aqd	>300	--	
M430	DDD-(BtA-Atmp) ₂	18	5.3	11.3
M432	DDD-(2Nal-Asp-Atmp) ₂	70	--	10
M436-1	TDIM-BtA-Atmp	8	4.8	
M436-2	TDIM-(BtA-Atmp) ₂	4.5	5.8	Ag
M438	F5c-3Pal-Atmp	26	4.8	10
M440	Dcg-BtA-Atmp	30	--	
M448	Dmac-BtA-Atmp	2.7	5.7	Ag
M456	F5c-Cys(Meb)-Atmp	4.7	5.3	Inact
M460	DDD-(3Pal-Nal-Cyh) ₂	15	Wk	
M466	F5c-2Nal-3Ampy	Inact	Wk	
M470	F5c-2Nal-Ampz	11	5.2	Inact
M472	Dmac-2Nal-Ampz	25	5.3	Inact
M474	Pya-2Nal-3Abza	35	Wk	Inact
M476	Tha-BtA-Atmp	15	4.6	Inact
M478	Dmac-2Nal-Thm	30	5.1	Inact
M480-1	HOOC-DDD-Pac-2Nal-Ampz	45	5.1	
M480-2	DDD-(Pac-2Nal-Ampz) ₂	--	5.6	
M492	F5c- <i>m</i> ABz-2Nal-Ampz	45	5.1	Inact

M506	Mse-Pac-BtA-Atmp	11	4.9	--
M526	F5c-2Nal-Dmp	10	5.4	Inact
M536	F5c-2Nal-Dmab	4	Wk	Inact
M538	DDD-(Pac-2-Nal-Dmp) ₂	>80	Wk	
M568	F5po-2Nal-Atmp	10	5.8	Ag
M570	F5c-OC2Y-Atmp	1.8	5.6	Ag
M572	Dca-2Nal-Acep	2.6	Wk	Wk
M574	Dns-Tyr(Bzl)Atmp	4.5	--	Inact
M582	Dmac-OC2Y-Atmp	3	5.4	9.5
M584-A	DDD-[DArg(Tos)-2Nal-Atmp] ₂	5	Inact	10.3
M584-B	DDD-(DArg-2Nal-Atmp) ₂	5	5.7	11.3
M586-A	Mse-Pac-Igl-Atmp	15	5.3	12
M586-B	Seb-Pac-Igl-Atmp	40	Wk	12.3
M588	α-DDD-(Lys-DArg-2Nal-Atmp) ₂	9.4	Wk	10
M592	F5c-OC2Y-Matp	1.5	4.9	Ag
M594	F5c-MC2Y-Atmp	3.7	5.0	8
M594	F5c-MC2Y-Atmp	3.7	5.0	8
M596-A	DDD-[Arg(Tos)-2Nal-Atmp] ₂	15	5.0	
M596-B	DDD-(Arg-2Nal-Atmp) ₂	8.2	Wk	Inact
M602	Chc-OC2Y-Atmp	12	--	10.8
M604	Pac-2Nal-Ecap	43	4.5	
M606	DDD-(Pac-2Nal-Api) ₂	30	5.0	10
M614	F5c-(N-Dmb)-Tyr(Bzl)-OMe	9.1	Wk	
M616	DDD-(Pac-1Nal-Atmp) ₂	1.4	5.4	--
M618	F5c-DArg-2Nal-Arg-Matp	18	--	--
M620	DDD-(DArg-2Nal-Arg-Matp) ₂	2.0	5.5	--
M622	F5c-OC2Y-Mapp	1.2	5.7	--
M624	Dns-OC2Y-Matp	1.4	5.1	--
M626	Pya-OC2Y-Matp	3.7	4.8	--
M628	Cin-OC2Y-Matp	1.6	5.2	--
M630	Dmac-OC2Y-Matp	1.6	5.0	--
M632	Atcp-OC2Y-Matp	1.4	5.4	--
M636	DDD-(DArg-Arg-Aud-Pac-2Nal-Atmp) ₂	1.7	5.8	--
M638	DDD-(DArg-Igl-Arg-Matp) ₂	0.6	Inact	
M640	DDD-(DArg-BtA-Arg-Matp) ₂	3.0	5.9	
M648	F5c-PaF(Mes)-Atmp	Inact	5.0	
M650	Atcp-OC2Y-Mapp	3.7	--	--
M652	Ppa-OC2Y-Mapp	7.5	5.7	--
M654	Sul-Atmp	Inact	4.5	
M656	Sul-2Nal-Atmp	13	5.4	
M660	DDD-(His-1Nal-Atmp) ₂	30	Wk	--
M662	F5c-tLeu-Atmp	Inact	5.2	--
M664	F5c-OCIY-Matp	1.2	5.0	
M666	Dns-OCIY-Matp	1.3	5.0	
M668	SBEC-(DArg-2Nal-Arg-Matp) ₂	3.4	5.2	
M670	DTP-(DArg-Igl-Arg-Matp) ₂	Inact	5.1	
M672	HDD-(DArg-Igl-Arg-Matp) ₂	--	--	
M674	DDD-(DArg-F5F-Arg-Matp) ₂	3.5	Wk	
M678	(Dns-DArg-Igl-Arg) ₂ -DDA	1.1	5.3	
M724	F5c-DArg-Aud-OC2Y-Gly-Atmp	12	5.4	
M744	DDD-(DArg-2Nal-Arg-Dmab) ₂	3.4	5.3	
M798	F5c-OC2Y-Dmab	37	--	
M800	DDD-(DArg-OC2Y-Dmab) ₂	27	5.3	
M832	F5c-PFF-Dmab	47	4.6	
M834	DDD-(DArg-PFF-Arg-Dpea) ₂	1.6	5.3	
M848	DDD-(DArg-F5F-Arg-Dmab) ₂	--	--	

M880	DDD-(DArg-F5F-Arg-Dpea) ₂	--	--
M886-1	DDD-DArg-PFF-Arg-NH ₂ L-DArg-PFF-Arg-Dpma	3.2	Wk
M886-2	DDD-(DArg-PFF-Arg-Dpma) ₂	--	Inact
M892	DDD-(DArg-PFF-Arg-PFF-NH ₂) ₂	8.5	Wk
M900	DDD-(DArg-F5F-Arg-PaF-NH ₂) ₂	6.3	--
M916	F5c-DArg-PFF-Arg-PFF-NH ₂	5.7	4.9
M1032	DDD-(DArg-Igl-Mapp) ₂	15	5.4
M1040	EDTA-(OC2Y-Atmp) ₄	0.73	--

5

Footnotes:

^aED₅₀ for killing of SCLC strain SHP-77 *in vitro*, μM.

^bpA₂ for bradykinin antagonist activity on isolated guinea pig ileum. The pD₂ of

10 bradykinin is 7.4 on ileum. Higher numbers indicate higher potency.

^cpA₂ for bradykinin antagonist potency on cloned human B2 receptors, pM. The pD₂ for bradykinin is 11. Higher numbers indicate higher potency.

Ag = agonist; Inact = inactive; Wk = weak

5 Table 3. ACTIVITIES OF CYCLIC PEPTIDES

NUMBER	STRUCTURE	MTT ^a	GPI ^b
B9458-2	DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-CpGΨ(CH ₂ N)Arg _____CO_____CH ₂ _____/ \	--	6.1
	CH ₂ CO ₂ H		
B9462	Darg-Arg-Pro-Hyp-Gly-Thi-Ser-Digl-CpGΨ(CH ₂ N)Arg _____CO_____CH ₂ _____/ \	7.3	6.0
	CH ₂ CO ₂ H		
B10302	c[DArg-Arg-Pro-Hyp-Gly-Igl-Ser-Dlgl-Oic-Arg]	Inact	5.2
B10304	Aca-c[DArg-Arg-Pro-Hyp-Gly-Thi-Ser-Nig-Oic-Arg]	Inact	6.4
B10306	c[Arg-DNMF-DTrp-Phe-DTrp-Leu]	Inact	Wk
B10312	α-DDD-(c[Lys-DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DF5F-Oic-Arg]) ₂	3.8	--
M680	c[DArg-Arg-Eac-Ser-DF5F-Nc7G-Arg]	--	--
M824	c[Ava-Igl-Ser-DF5F-Oic-Arg]	Inact	5.2
M850	c[DArg-Arg-Aud-Dlgl-PFF-Arg]	1.4	5.1
M868-2	c[DArg-Arg-Aud-DF5F-Oic-Arg]	9.2	6.1
M870	c[DArg-Arg-Add-DF5F-Oic-Arg]	5.5	5.3
M872	c[DArg-Arg-Eac-Ser-DF5F-Oic-Arg]	2.2	Inact
M874	c[DArg-Arg-Add-Ser-DF5F-Oic-Arg]	11	5.0
M876	c[DArg-Arg-Aud-Ser-DF5F-Oic-Arg]	22.5	5.4
M878	c[DArg-Arg-Add-Dlgl-PFF-Arg]	7	Wk
M882	c[DArg-Arg-Add-Ser-Dlgl-PFF-Arg]	4.5	Inact
M896	c[DArg-Arg-Eac-Dlgl-PFF-Arg]	65	Wk
M902	c[DArg-Arg-Ava-Ser-Dlgl-PFF-Arg]	30	5.5
M906	c[DArg-Arg-Eac-DF5F-Oic-Arg]	45	Wk
M908	c[DArg-Arg-Ava-Ser-DF5F-Oic-Arg]	40	4.9
M910	c[Bala-DArg-Arg-Eac-Ser-DF5F-Oic-Arg]	42	5.2
M924	c[Suc-DArg-Arg-Eac-Ser-Dlgl-PaF]-Arg	37	Wk
M934	c[DNiK-Arg-Eac-Ser-DF5F-Oic-Arg]	--	--
M940	c[DNiK-PzO-Eac-Ser-DF5F-Oic-Arg]	--	--
M986	c[Add-DArg-F ₅ F-Arg]	--	--

Footnotes:

10 ^aED₅₀ for killing of SCLC strain SHP-77 *in vitro*, μM.

^bpA₂ for bradykinin antagonist activity on isolated guinea pig ileum. The pD₂ of bradykinin is 7.4 on ileum. Higher numbers indicate higher potency.

Inact = inactive; Wk = weak

5 **Table 4. PREVIOUSLY DESCRIBED KNOWN PEPTIDES THAT NEWLY SHOW
IN VIVO ANTI-CANCER ACTIVITY**

NUMBER	STRUCTURE
B9430	DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg
B9330	DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Nig-Arg
B10044	DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg
B10050	Lys-Lys-Arg-Pro-Hyp-Gly-Igl-Ser-DTic-ChG
B10206	DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Nc7G-Arg
B10288	DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Oic-Arg

10 These compounds showed anti-tumor activity *in vivo* when tested by the
procedure of Example XXI.

**TABLE 5. CYTOTOXICITY IN VITRO AGAINST STRAINS OF PROSTATE
CANCER**

COMPOUND NUMBER	PROSTATE CANCER CELL LINE					SCLC
	DU14	TSU	LNCa	PC-3	PPC1	
B9870	0.08	6.5	3.7	3.2	4.3	0.15
M570	1.2	2.8	3.0	1.6	3.0	1.8
M590	0.01	7.0	7.0	6.3	12	4.5

15

Numbers are ED₅₀ (μM) for cytotoxic activity. Activity against SCLC strain SHP-77 is included for comparison.

TABLE 6. ABBREVIATIONS USED FOR COMPOUNDS

B9430 = DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg
 B9870 = SUIM-(DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg)₂
 Aaa = 1-Adamantaneacetyl
 AAA = amino acid analysis
 ABz = Aminobenzoic acid
 ABza = Aminobenzyl alcohol
 Ac = Acetyl
 Ac3c = 1-Amino-1-cyclopropanecarboxylic acid
 Ac5c = 1-Amino-1-cyclopentanecarboxylic acid ("cyclo-Leu")
 Ac6c = 1-Aminocyclohexanecarboxylic acid
 Aca = 1-Adamantanecarboxyl
 Acep = 4-Amino-1-carbethoxymethyl-2,2,6,6-tetramethylpiperidine
 ADA = 1,3-Adamantanediacyl
 Add = 12-Aminododecanoic acid
 Aib = α-Aminoisobutyric acid
 Aic = 2-Aminoindane-2-carboxylic acid

AIG = α -Allylglycine (2-amino-4-pentenoic acid)
 Ama = Aminomethylanthranilic acid
 Amb = Aminomethylbenzoic acid
 Ampy = 3-Aminomethylpyridine
 Ampz = 1-Amino-4-methylpiperazine
 Ana = Anthranilic acid
 APa = *p*-Aminophenylacetic acid
 APb = *p*-Aminophenylbutyric acid
 ApC = S-3-Aminopropylcysteine
 Api = 4-Aminopiperidine
 Apmp = 4-Amino-1,2,2,6,6-pentamethylpiperidine
 Atp = 4-Amino-1-phenylmethyl-2,2,6,6-tetramethylpiperidine
 Aqd = 4-Aminoquinaldine
 Aqu = 3-Aminoquinuclidine
 Arg(NO₂) = Arginine(Nitro)
 Atc = 2-Aminotetralin-2-carboxylic acid
 Atcp = 4-Amino-3,5,6-trichloropicolinic acid
 Atmp = 4-Amino-2,2,6,6-tetramethylpiperidine
 AtmpO = 4-Amino-2,2,6,6-tetramethylpiperidinyloxy
 Atpc = 4-Amino-2,2,6,6-tetramethyl-4-piperidinecarboxylic acid
 Atpm = 4-Amino-4-methoxycarbonyl-2,2,6,6-Tetramethylpiperidine
 (4-Amino-2,2,6,6-tetramethyl-4-piperidinecarboxylic acid methyl
 ester)
 Aud = 11-Aminoundecanoic acid
 Ava = 5-Aminovaleric acid
 Azt = Azetidine-2-carboxylic acid
 BAla = β -Alanine
 BApG = N,N-bis(3-aminopropyl)-glycine
 BAPTA = 1,2-bis(2-Aminophenoxy)ethane-N,N,N',N'-tetraacetyl
 Bip = Biphenylalanine
 Boc = (*tert*-Butoxycarbonyl); [(1,1-dimethylethoxy)carbonyl]
 BOP = Benzotriazoyloxytris(dimethylamino)phosphonium
 hexafluorophosphate
 BPHD = N,N'-bis(2,2,6,6-tetramethyl-4-piperidiny)-1,6-hexanediamine
 BSH = 1,6-Bissuccinimidohexane
 BtA = 3-Benzothienylalanine
 BTAC = Benzene-1,3,5-*tris*-carboxamido-6-caproyl
 BTC = 1,3,5-Benzenetricarboxyl
 Bz = Benzoyl
 Bzl = Benzyl
 CAch = *cis*-2-Amino-1-cyclohexanecarboxylic acid
 Cca: 2-Chlorocinnamic acid
 CDF = *p*-Chloro-D-phenylalanine
 ChA = α -Cyclohexylalanine
 Chc = α -Cyano-4-hydroxycinnamoyl

ChG = α -Cyclohexylglycine
CHO = Chinese hamster ovary
CHTC = 1,3,5-Cyclohexanetricarboxyl
CHyp = *cis*-4-Hydroxy-proline
Cin = Cinnamoyl
CMeb = S-(4-Methylbenzyl) cysteine
CmF = (Z) *p*-Chloro-2,3-methanophenylalanine
Cmp = 4-Carboxymethylpiperazine
CpA = α -Cyclopropylalanine
CpG = α -Cyclopentylglycine
CpG Ψ (CH₂N)Arg = CpG pseudo(CH₂NH) Arg
CPTA = *trans*-1,2-Diaminocyclohexane- N,N,N',N'-tetraacetyl
CTAC = Cyclohexane-1,3,5-*tris*-carbamido- ϵ -caproyl
Ctim = 13-Carboxytridecanimidyl
Cyh = Cyclohexylamine
Dabz = Diaminobenzoic acid
DArg(NO₂) = Nitro-Arginine
Dca = Dicyclohexylacetyl
Dcg = N,N'-Dicyclohexylguanidyl
DCM = Dichloromethane
DDA = 1,10-Decanediamine
DDD = Dodecanedioyl
DDS = 2-Dodecen-1-ylsuccinyl
DEA = N,N'-Diethylethylenediamine
DhP = 3,4-Dehydroproline
Dhq = 2,3-Dehydroquinuclidine-3-carboxyl
DIC = Decahydroisoquinoline-3-carboxylic acid
DIEA = Diisopropylethylamine
Dmab = 4-Dimethylaminobenzylamine
Dmac = 4-Dimethylaminocinnamyoyl
Dmb = 4-(Dimethylamino)benzyl
DmF = 2,4-Dimethylphenylalanine
DMF = Dimethyl formamide
DmK = ϵ -Dimethyllysine
Dmm = 2,6-Dimethylmorpholine
Dmp = 3-Dimethylaminopropylamine
DmtP = 5,5-Dimethyl-4-thiaproline
Dns = Dansyl (5-dimethylamino-1-naphthalenesulfonyl)
22Dp = 2,2-Diphenylpropionyl
33Dp = 3,3-Diphenylpropionyl
Dpa = Diphenylacetyl
Dpea = Diphenylethylamine
Dpma = Diphenylmethylamine
Dpr = 2,3-Diaminopropionic acid
DTP = Dithiobis-propionyl

DTPA = Diethylenetriaminepentaacetyl
Eac = ϵ -Aminocaproic acid
Ecap = N-Ethoxycarbonyl-4-amino-piperidine (Ethyl 4-amino-1-piperidinecarboxylate)
EDA = 4,4'-Ethylenedianiline
EDP = 4,4'-Ethylenedipiperidine
EDTA = Ethylenediaminetetraacetyl
EDTP = Ethylenediaminetetrapropionic acid
EGS = Ethylene glycol-*bis*-succinyl
EGTA = Ethylene glycol-*bis*(β -aminoethyl ether)- N,N,N',N'-tetraacetyl
EOPC = 1,1'-Ethylenebis(5-oxo-3-pyrrolidinecarboxyl)
ETTA = 2,2',2'',2'''-[Ethanediylidenetetrakis(thio)tetrakisacetyl
F2F = Difluorophenylalanine
F5bz = Pentafluorobenzoyl
F5c = 2,3,4,5,6-Pentafluorocinnamoyl
F5F = Pentafluorophenylalanine
F5pa = 2,3,4,5,6-Pentafluorophenylacetyl
F5po = 2,3,4,5,6-Pentafluorophenoxyacetyl
Fbz = *para*-Fluorobenzoyl
Flu = Fluorescein thiourea
Gaa = Guanidinoacetyl
GaP = 2-Guanidyl-3-(4-aminophenyl)propionic acid
Gbz = 4-Guanidinobenzoyl
Glt = Glutaryl
Gun = Guanidyl
HATU = O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
HbQ = 4-Hydroxybutylglutamine
HDA = 1,6-Hexanediamine
HDD = Hexadecanedioyl
HF = Hydrogen fluoride
HFG = Hexafluoroglutaroyl
HiG = Hexahydro-2-indanylglycine
HOAt = 1-Hydroxy-7-azabenzotriazole
hPhe = Homo-phenylalanine
HPLC = high performance liquid chromatography
Hxa = Hexanoic acid
Hyp = *trans*-4-Hydroxyproline
Ica = Indoline-2-carboxylic acid
Igl = α -2-Indanylglycine
Ing = α -1-Indanylglycine
Inp = Isonipectic acid
Iq2a = 6,7-Dimethoxy-3,4-dihydro-1-isoquinolineacetic Acid
Iq4a = 6,7-Dimethoxy-1,2,3,4-tetrahydro-1-isoquinolineacetic acid
Lau = Lauroyl


Leu(r)Leu = Leu-pseudo(CH₂NH)Leu
LeuΨ(CH₂NH)Leu = Leu-pseudo(CH₂NH)Leu
LDMS = laser desorption mass spectrometry
mA₂Bz = 3,5-Diaminobenzoic acid
MaG = α-Methallylglycine (2-amino-3-methyl-4-pentenoic acid)
Mapp = 4-(Methylamino)-1,2,2,6,6-pentamethylpiperidine
Matp = 4-(Methylamino)-2,2,6,6-tetramethylpiperidine
MatpO = 4-(N-methylamino)-2,2,6,6-tetramethylpiperidinyloxy
MBC = S-(4-methylbenzylcysteine
MBHA = Methylbenzhydrylamine
MC2Y = N-Methyl-O-2,6-dichlorobenzyl-tyrosine
Mca = 2-Methylcinnamic acid
Mcg = Monocyclohexylguanidyl
Meb = Methylbenzyl
MeP = 2,4-Methanoproline
Mes = Methanesulfonyl
MFE = (E)-2,3-Methanophenylalanine
MFF = *meta*-Fluorophenylalanine
Mosi = Methoxy-suberimido
Moti = 14-Methoxytetradecanediimidoyl
Mse = Methoxysebacyl
MTT = (3-(4,5)-Dimethyltriazol-2-yl)-2,5-diphenyl tetrazolium bromide
Nal = β-Naphthylalanine
Nap = Naphthoyl
Nba = Norbornane-2-acetyl
Nbc = Norbornenedicarboxyl
Nbi = Norbornenedicarboximide
Nbn = 2-Aminonorbornane-2-carboxylic acid
Nc5G = N-Cyclopentylglycine
Nc6G = N-Cyclohexylglycine
Nc7G = N-Cycloheptylglycine
Nc8G = N-Cyclooctylglycine
Nig = N-2-Indanylglycine
NiK = ε-Nicotinoyllysine
NMF = N-Methylphenylalanine
NSCLC = non-small cell carcinoma
OBS = O-Benzylserine
OBT = O-Benzylthreonine
OBY = O-Benzyltyrosine
OC2Y = O-2,6-Dichlorobenzyltyrosine
OCiY = O-2,6-Dichlorobenzyl-3,5-diiodotyrosine
Oct = Octanoyl
Oic = Octahydroindole-2-carboxylic acid
OMe = O-Methyl
OMY = O-Methyltyrosine

OSY = Tyrosine O-sulfate ester
Paa = Phenylacetyl
Pac = 4-Aminocinnamic acid
PaF = *p*-Aminophenylalanine
Pal = β -Pyridylalanine
Pba = Phenylbutyryl
Pcc = *trans*-2-Phenyl-1-cyclopropanecarboxylic acid
PCF = *p*-Chlorophenylalanine
Pcpa = α -Phenylcyclopentaneacetyl
PdF = *p*-Dicyclohexylguanidylphenylalanine
PFF = *p*-Fluorophenylalanine
PFS = Perfluorosuberoyl
PgF = *p*-Guanidinophenylalanine
PheOL = Phenylalaninol
PhG = Phenylglycine
Pip = Pipelic acid ("homo-Pro")
PipA = β -3-Piperidylalanine
PNF = *p*-Nitrophenylalanine
Ppa = Phenylpropionyl
Pya = *trans*-3-(3-Pyridyl)acryloyl
PyAOP = 7-Azabenzotriazol-1-yloxytris(pyrrolidino)phosphonium hexafluorophosphate
PzK = ϵ -Pyrazinoyllysine
PzO = 4-Pyrazinoylornithine
Saa = *trans*-Styrylacetic acid
SBEC = Sulfo-*bis*-ethoxycarbonyl
Sbl = Sebacoyl
SCLC = small cell lung carcinoma
Seb = Sebacyl
Sin = Sinapinyl (3,5-dimethoxy-4-hydroxycinnamoyl-)
Ste = Stearoyl
Sua = Sulfanilamide (4-Aminobenzenesulfonamide)
SUB = Suberyl
Suc = Succinyl
Sud = Sulfadiazine
SUIM = Suberimidyl
Sul = Sulindac
Tba = *t*-Butyl-acetyl
TDIM = Tetradecanediimidyl
TEA = Triethylamine
TFA = Trifluoroacetic acid
Tfmc = *trans*-4-(Trifluoromethyl)cinnamoyl
Tha = 3-(2-Thienyl)acryloyl
Thi = β -2-Thienylalanine
Thm = Thiomorpholine

Thz = Thiazolidine-4-carboxylic acid (4-thiaproline)
Tic = 1,2,3,4-Tetrahydroisoquinoline-3-carboxylic acid
TLC = thin layer chromatography
TLeu = *tert*-Leucine
TMF = 2,4,6-Trimethylphenylalanine
Tos = *p*-Toluenesulfonyl
Tpac = 2,2,5,5-Tetramethyl-3-(aminoethyl)-pyrroline-3-carboxamide
TREN = *tris*(2-Aminoethyl)amine
Trx = Tranexamic acid (*trans*-4-((Aminomethyl))cyclohexanecarboxylic acid)


5 WHAT IS CLAIMED IS:

1. A compound of the formula:



10

wherein X is a linker group having 2-5 functional groups or is absent, $n = 1$, and Formula II is



15

wherein R, A, B, C, D, E, F, G, H, I, J, and K are selected from the following or may be absent, and wherein K is Arg or an Arg derivative:

R	A -1	B 0	C 1	D 2	E 3	F 4	G 5	H 6	I 7	J 8	K 9
Absent or 3,3DP Aaa Ac	Absent or DmK Lys Lys(eF lu)	Absent or Apc Arg DArg	Absent or ApC Arg DmK	Absent Or MeP Nig NMF	Absent or Hyp Pro	Absent or Ava BAIa Dpr	Absent or Add Aud CpG	Absent or Arg Gly Pac	Absent or 2Nal DCpG DF5F	Absent or 2Nal 2Nal- NH ₂	Absent or Arg Arg(H) Arg- CH ₂ O H Arg- NH ₂ Arg(N O ₂) Arg- OMe DArg DArg- NH ₂ DArg(NO ₂)
Aca	NiK	DLys	NiK	Pro		Eac	DDM F	Pac	Dlgl	3,4F2 F	Arg- NH ₂ Arg(N O ₂) Arg- OMe DArg DArg- NH ₂ DArg(NO ₂)
BAPg	PzO	DmK	NiO			Gly	DMF	Ser	DPFF	3Pal	Arg(N O ₂) Arg- OMe DArg DArg- NH ₂ DArg(NO ₂)
Cca		DniK	PaF				Eac	Thr	DPhe	Ac6c	Arg- OMe DArg DArg- NH ₂ DArg(NO ₂)
Cin Dca		DpaF DPzK	PzO				Igl Lys		DTic Gly	Aic Ana	DArg DArg- NH ₂ DArg(NO ₂)
Dcg		DPzO					Pac		mABz	Apb	DArg(NO ₂)
Dhq Dmac Dpa		Lys NiK PaF					Phe Thi		pABz Pac PaF(D cg) pAmb	Apb Atpc Bip	DArg DArg- NH ₂ DArg(NO ₂)
F5bz F5c		PzO DArg- (NO ₂)	Arg- (NO ₂)							Cmp CpG	DArg DArg- NH ₂ DArg(NO ₂)
F5pa Gun										DhPhe Dpr(F bz) Dpr(Pa a)	DArg DArg- NH ₂ DArg(NO ₂)
Hxa										F5F F5F- NH ₂ Hphe lca Igl Igl-	DArg DArg- NH ₂ DArg(NO ₂)
Mca Mcg											DArg DArg- NH ₂ DArg(NO ₂)
Moti Pcc Ppa Pya											DArg DArg- NH ₂ DArg(NO ₂)

Saa
Ste


Tfmc

NH₂
Ileu
Lys(C
H₃)₃
Lys(F
5bz)
Mapa
MBC
MFF
Nc6G
Nc7G
NMF
OBS
OBT
OBY
OC2Y
Oic
Oic-
NH₂
PABz
Pac
PaF(F
5c)
PaF(F
bz)
PaF(M
cg)
PaF(P
pa)
PaF(Si
n)
*p*Amb
*p*APa
PCF
PdF
PFF
PFF-
NH₂
Phe
PNF
Thi
Tic
Trp
Trx
Tyr

- 5 2. A compound of the formula:



10 wherein X is a linker group having 2-5 functional groups or is absent, n = 1, and Formula III is




15 wherein R, A, B, C, D, E, and F are selected from the following or may be absent, and wherein F is not Arg or an Arg derivative:

R	A 1	B 2	C 3	D 4	E 5	F 6
Absent or	Absent or	Absent or	Absent or	Absent or	Absent or	Absent or
2,2Dp 3,3Dp	DArg DArg(NO ₂)	Arg	Add Aud	2Nal 3Pal	1Nal 2Nal	2Nal 3Pal
Aaa Ac Aca Boc Chc Cin			Ava Eac Lys Pac	Arg Arg(Tos) Atcp D2Nal DArg DArg(Tos)	2Nap 3Pal Apa Arg Arg-NH ₂ Asp	ABza ABza Ama Ampy Ampz Apa
Ctim Dca Dcg Dhq Dmac Dns				DF5F DIgl DPFF Eac F5F Gly	Atc Atcp Bip BtA Cys(Meb) Cys(SO ₃ H)	Api Aptp Aqd Aqu Arg(H) Arg-CH ₂ OH
Dpa F5c F5pa F5po Gbc Gun Hxa Mcg Mse Pya Seb Sin				His Igl mABz OC2Y Pac PFF	D2Nal DArg DArg-NH ₂ F5F Glu Gly Igl Inp Iqa mABz MC2Y N-Dmb- Tyr(Bz)- OMe OC2Y OCIY Oic pABz	Arg-NH ₂ Arg-OMe Asp Asp(Aqu) Atcp Atmp AtmpO Atpm Cyh Dmab Dmm Dmp
Sul Tfmc Tha						Dpea Dpma Dpr(Dcg-2-Nap) Ecap

PaF(Mes)	F5F-NH ₂
PFF	GaP
Tic	<i>m</i> A ₂ Bz
<i>t</i> Leu	<i>m</i> A ₂ Bz(Dc g)
Trp	<i>m</i> A ₂ Bz(Gu n)
Try	<i>m</i> ABz
Try(Bzl)	Mapp
Tyr	Matp
Arg(NO ₂)	MatpO
	<i>p</i> ABz
	PaF
	PaF(Dcg)
	PaF(Mcg)
	PaF-NH ₂
	PFF-NH ₂
	PgF
	PzO
	Sud
	Thm
	Thm
	Tpac
	Tpac
	Tyr(Bz)O
	Me

5

3. A compound of the formula:

10 wherein X is a linker group having 2-5 functional groups or is absent, $n = 1$, and Formula IV is

15


wherein A, B, C, D, E, F, G, H, I, J, K and L are selected from the following or may be absent:

A	B	C	D	E	F	G	H	I	J	K	L
0	1	2	3	4	5	6	7	8	9	10	11
Absent	Absent	Absent	Absent	Absent	Absent	Absent	Absent	Absent	Absent	Absent	Absent
or	or	or	or	or	or	or	or	or	or	or	or
DArg	Arg	Pro	Lys	Pro	DTrp	Gln	DTrp	Phe	DTrp	Leu(r)	Leu-NH ₂
	DArg					DNM					Leu
						F					

- 5 4. A compound of the formula:




- 10 wherein X is a linker group having 2-5 functional groups or is absent; $n = 1$; c indicates cyclization, the site of cyclization selected from the group consisting of the c-terminus, and a side chain functional group; and Formula V is



- 15 wherein X, A, B, C, D, E, F, G, H, I, J, and K are selected from the following or may be absent:

X	A	B	C	D	E	F	G	H	I	J	K
	-1	0	1	2	3	4	5	6	7	8	9
Absent or α -Aca 3,3Dp	Absent or Ava BAIa DmK Glt Lys Suc	Absent or DArg DNik DPaF DPzK DPzO	Absent or Arg NiK PzO	Absent or Pro	Absent or Hyp	Absent or Gly	Absent or Add Aud Ava BAIa DNMF Eac Igl Thi	Absent or DArg Ser Thr	Absent or DDab DDpr DF5F DIgl DLys DOm DPaF Nig Pac Phe	Absent or DTrip F5F Lys Nc7G Oic PaF PFF Phe	Absent or Arg Leu NiK PaF 3Pal

- 20 5. A compound of the formula



wherein [ACA] is a compound of claim 1, 2, 3, or 4.

- 25 6. A method to inhibit tumor growth in an animal in need of such inhibition, comprising administering a compound selected from the group consisting of a compound of claim 1,
 30 DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg,
 DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Nig-Arg,
 DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg,
 Lys-Lys-Arg-Pro-Hyp-Gly-Igl-Ser-DTic-ChG,
 DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Nc7G-Arg, and
 DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Oic-Arg.

5

7. A method to inhibit tumor growth in an animal in need of such inhibition, comprising administering a compound selected from the group consisting of a compound of claim 2,

- 10 DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg,
DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Nig-Arg,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg,
Lys-Lys-Arg-Pro-Hyp-Gly-Igl-Ser-DTic-ChG,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Nc7G-Arg, and
DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Oic-Arg.

15

8. A method to inhibit tumor growth in an animal in need of such inhibition, comprising administering a compound selected from the group consisting of a compound of claim 3,

- 20 DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg,
DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Nig-Arg,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg,
Lys-Lys-Arg-Pro-Hyp-Gly-Igl-Ser-DTic-ChG,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Nc7G-Arg, and
25 DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Oic-Arg.

9. A method to inhibit tumor growth in an animal in need of such inhibition, comprising administering a compound selected from the group consisting of a compound of claim 4,

- 30 DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg,
DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Nig-Arg,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg,
Lys-Lys-Arg-Pro-Hyp-Gly-Igl-Ser-DTic-ChG,
35 DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Nc7G-Arg, and
DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Oic-Arg.

10. A method to inhibit tumor growth in an animal in need of such inhibition, comprising administering a compound selected from the group consisting of a compound of claim 5, DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg,

- 40 DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Nig-Arg,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg,
Lys-Lys-Arg-Pro-Hyp-Gly-Igl-Ser-DTic-ChG,
45 DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Nc7G-Arg, and
DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Oic-Arg.

- 5 11. A method to induce apoptosis, comprising administering a compound selected from the group consisting of a compound of claim 1,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg,
DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Nig-Arg,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg,
10 Lys-Lys-Arg-Pro-Hyp-Gly-Igl-Ser-DTic-ChG,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Nc7G-Arg, and
DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Oic-Arg.
12. A method to induce apoptosis, comprising administering a compound selected from
15 the group consisting of a compound of claim 2,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg,
DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Nig-Arg,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg,
Lys-Lys-Arg-Pro-Hyp-Gly-Igl-Ser-DTic-ChG,
20 DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Nc7G-Arg, and
DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Oic-Arg.
13. A method to induce apoptosis, comprising administering a compound selected from
the group consisting of a compound of claim 3,
25 DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg,
DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Nig-Arg,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg,
Lys-Lys-Arg-Pro-Hyp-Gly-Igl-Ser-DTic-ChG,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Nc7G-Arg, and
30 DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Oic-Arg.
14. A method to induce apoptosis, comprising administering a compound selected from
the group consisting of a compound of claim 4,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg,
35 DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Nig-Arg,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg,
Lys-Lys-Arg-Pro-Hyp-Gly-Igl-Ser-DTic-ChG,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Nc7G-Arg, and
DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Oic-Arg.
40
15. A method to induce apoptosis, comprising administering a compound selected from
the group consisting of a compound of claim 5,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DIgl-Oic-Arg,
DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Nig-Arg,
45 DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Oic-Arg,
Lys-Lys-Arg-Pro-Hyp-Gly-Igl-Ser-DTic-ChG,
DArg-Arg-Pro-Hyp-Gly-Igl-Ser-DF5F-Nc7G-Arg, and
DArg-Arg-Pro-Hyp-Gly-Thi-Ser-DTic-Oic-Arg.

Fig. 1 INHIBITION OF GROWTH OF SCLC SHP-77 BY B10054 *in vivo*
Daily IP Injection




Fig. 2 INHIBITION OF GROWTH OF NSCLC A-549 BY M620 *in vivo*




Fig. 3 INHIBITION OF GROWTH OF SCLC SHP-77 BY B9430 *in vivo*
Bradykinin antagonists/SHP77/nude mice
5 mg/kg daily IP injection




Fig. 4 INHIBITION OF GROWTH OF SCLC SHP-77 BY B10238 *in vivo*




Fig. 5 INHIBITION OF GROWTH OF SCLC SHP-77 BY M570/TFA

And M570/HCl *in vivo*

SHP77/nude mice/BKM analogs

5 mg/kg daily IP injection




Fig. 6 INHIBITION OF GROWTH OF SCLC SHP-77 BY M822 *in vivo*





Fig. 7 INHIBITION OF GROWTH OF SCLC SHP-77 BY M638 *in vivo*



INTERNATIONAL SEARCH REPORT

 International application No.
PCT/US99/19381
A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) :C07K 7/06, 7/18, 14/00, 19/00, 99/18; A61K 38/00, 38/08

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/13, 15; 530/314, 326, 327, 328, 402, 408, 807, 815, 816; 548/314.4, 521; 435/107, 117, 118, 121, 128, 129

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

File CA; File HCAPLUS; File USPATFULL

Structures and Inventors' name searched

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	US 5,635,593 A (CHERONIS et al) 03 June 1997, the entire document, especially col. 7, line 45; col. 8, lines 32-49; col. 24, TABLE B and abstract.	1 ----- 2-5
X --- Y	WO 97/09347 A1 (CORCETH INC) 13 March 1997, especially TABLE I and claims.	1, 6 ----- 2-5, 7-15
Y	CHAN, D. et al Novel Bradykinin Antagonist Dimers for the Treatment of Human Lung Cancers. Immunopharmacology. 1996, Vol. 33, pages 201-204, especially page 202 and Fig 1.	1-15

☒ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

31 JANUARY 2000

Date of mailing of the international search report

10 FEB 2000

 Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

FATEMEH T. MOEZIE

Telephone No. (703) 308-0193

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/19381

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X --- Y	GERA, L. et al. New Bradykinin Antagonists Having High Potency at Both B1 and B2 Receptors. Peptides: Chemistry, Structure and Biology Pravin. T.P. Kaumaya and Robert S. Hodges (Eds.) Mayflower Scientific Ltd., 1996, especially Table 1.	1 ----- 2-5

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/19381

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

514/13, 15; 530/314, 326, 327, 328, 402, 408, 807, 815, 816; 548/314.4, 521; 435/107, 117, 118, 121, 128, 129

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record.**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.